

ELSEVIER

Mechanisms of Development 55 (1995) 235



Volume contents

Volume 55 (1996)

Issue contents	1	K. Kostakopoulou, A. Vogel, P. Brickell, C. Tickle 'Regeneration' of wing bud stumps of chick embryos	
J. Joore, C. Fasciana, J.E. Speksnijder, W. Kruijer, O.H.J. Destrée, A.J.M. van den Eijnden-van Raaij,		and reactivation of Msx-1 and Shh expression in response to FGF-4 and ridge signals	119
S.W. de Laat, D. Zivkovic		aw	
Regulation of the zebrafish goosecoid promoter by mesoderm inducing factors and Xwnt1	3	C. Mason, M. Lake, A. Nebreda, R. Old A novel MAP kinase phosphatase is localised in the branchial arch region and tail tip of Xenopus embryos	
T. Hollemann, R. Schuh, T. Pieler, R. Stick Xenopus Xsal-1, a vertebrate homolog of the region		and is inducible by retinoic acid	133
specific homeotic gene spalt of Drosophila	19	B. Kablar, R. Vignali, L. Menotti, M. Pannese, M. Andreazzoli, C. Polo, M.G. Giribaldi, E. Boncinelli, G.	
J. Luo, H.M. Sucov, JA. Bader, R.M. Evans, V.		Barsacchi	
Giguère		Xotx genes in the developing brain of Xenopus laevis	145
Compound mutants for retinoic acid receptor (RAR) β			
and RAR α 1 reveal developmental functions for multiple RAR β isoforms	33	E. Sánchez-Herrero, J.P. Couso, J. Capdevila, I. Guerrero	
ipio ti ity isolomi		The fu gene discriminates between pathways to con-	
S.L. Helfand, B. Naprta		trol dpp expression in Drosophila imaginal discs	159
The expression of a reporter protein, β -galactosidase,		and the conference of the conf	
is preserved during maturation and aging in some cells		O. Hobert, I. Sures, T. Ciossek, M. Fuchs, A. Ullrich	
of the adult Drosophila melanogaster	45	Isolation and developmental expression analysis of Enx-1, a novel mouse Polycomb group gene	171
T. Bossing, G.M. Technau, C.Q. Doe			
huckebein is required for glial development and axon		K.M. Catron, H. Wang, G. Hu, M.M. Shen, C. Abate-	
pathfinding in the neuroblast 1-1 and neuroblast 2-2		Shen	
lineages in the Drosophila central nervous system	53	Comparison of MSX-1 and MSX-2 suggests a mo- lecular basis for functional redundancy	185
S. Nef, I. Allaman, H. Fiumelli, E. De Castro, P. Nef			
Olfaction in birds: differential embryonic expression		S.M. Shimeld, I.J. McKay, P.T. Sharpe	
of nine putative odorant receptor genes in the avian		The murine homeobox gene Msx-3 shows highly re-	
olfactory system	65	stricted expression in the developing neural tube	201
V. Wilson, R.S.P. Beddington		J.C. Corbo, M. Levine	
Cell fate and morphogenetic movement in the late		Characterization of an immunodeficiency mutant in	
mouse primitive streak	79	Drosophila	211
S. Stein, R. Fritsch, L. Lemaire, M. Kessel		K.H. Kaestner, G. Schütz, A.P. Monaghan	
Checklist: Vertebrate homeobox genes	91	Expression of the winged helix genes fkh-4 and fkh-5 defines domains in the central nervous system	22:
Issue contents	131		
		Author index	23
L. Chatelin, M. Volovitch, A.H. Joliot, F. Perez, A.		Subject index	233
Prochiantz		Volume contents	235
Transcription factor Hoxa-5 is taken up by cells in		· · · · · · · · · · · · · · · · · · ·	
culture and conveved to their nuclei	111		